


CLAIMS

I claim:

1. A platinum(II) complex selected from the group consisting of compounds of the general formulae I to IV and physiologically acceptable addition salts of them,

5



wherein the radicals

10

R^1 and $R^{1'}$ are selected independently of one another from the group consisting of substituted or unsubstituted alkylene and alkenylene radicals, which can be substituted by halogen, alkyl, alkenyl, cycloalkyl, cycloalkenyl, aryl, hydroxy, carboxy, sulphate, phosphate and / or heterocyclic compounds,

15

R^2 , R^3 , $R^{2'}$, $R^{3'}$ are selected independently of one another from the group consisting of $-(CH_2)_m-OH$, $-H$, substituted or unsubstituted alkyl radicals, saturated and unsaturated cyclic radicals and heterocyclic compounds which can be substituted by halogen, hydroxy, carboxy, sulphate and / or phosphate,

m signifies a natural number from 2 to 5,

20

R^4 is selected from the group consisting of substituted or unsubstituted alkylene, alkenylene, cycloalkylene and cycloalkenylene radicals, and aromatic and heterocyclic radicals,

$X, X' = -S(O_2)-$,

$k, i = 0$ or 1 ,

$Y = -OH, -SO_3H$,


$n = 0$ or 1 , and

25

a is selected from the group consisting of halogenides, OH^- , OH_2 , carboxylate, sulphate and sulphonate

and mixtures of the compounds for use as prophylactic and / or therapeutic agent for the treatment of diseases.

2. A platinum(IV) complex with a single or double intramolecular cyclization selected from the group consisting of compounds of the general formulae Ia to IVa and physiologically acceptable addition salts of them,



wherein the radicals

R¹ and R^{1'} are selected independently of one another from the group consisting of substituted or unsubstituted alkylene and alkenylene radicals, which can be substituted by halogen, alkyl, alkenyl, cycloalkyl, cycloalkenyl, aryl, hydroxy, carboxy, sulphate, phosphate and / or heterocyclic compounds,

R², R³, R^{2'}, R^{3'} are selected independently of one another from the group consisting of -(CH₂)_m-OH, -H, substituted or unsubstituted alkyl radicals, saturated and unsaturated cyclic radicals or heterocyclic compounds which can be substituted by halogen, hydroxy, carboxy, sulphate and / or phosphate,

m signifies a natural number from 2 to 5,

R⁴ is selected from the group consisting of substituted or unsubstituted alkylene, alkenylene, cycloalkylene and cycloalkenylene radicals, and aromatic and heterocyclic radicals,

X, X' = -S(O₂)-,

k, i = 0 or 1,

Y = -OH, -SO₃H,

n = 0 or 1, and

a, c, c' are selected independently of one another from the group consisting of halogenides, OH⁻, OH₂, carboxylate, sulphate and sulphonate

and mixtures of the compounds for use as prophylactic and / or therapeutic agent in the treatment of diseases.

3. The platinum(II) complex according to Claim 1, wherein

$R^2, R^3, R^{2'}, R^{3'}$ are selected independently of one another from the group consisting of hydrogen, methyl and ethyl radicals.

4. The platinum(II) complex according to claim 1, wherein R^1 and $R^{1'}$ are selected

5 independently of one another from the group consisting of C_{2-5} -alkylene radicals and C_{2-5} -alkenylene radicals.

5. The platinum(II) complex according to claim 1, wherein R^4 is ethylene.

6. The platinum(II) complex according to claim 1, wherein the radicals have the following significance:

10 $R^1, R^{1'}$ = substituted or non-substituted alkylene,
 $k, i = 0$, and / or
 $Y = -OH$.

7. The platinum(II) complex according to claim 1, wherein the radicals have the following significance:

15 $R^1, R^{1'}$ = ethylene,
 $k, i = 0$,
 $Y = -OH$, and / or
 $n = 0$.

8. Use of at least one platinum(II) complex according to claim 1, for the manufacture of
20 a prophylactic and / or therapeutic agent for the treatment of tumor diseases.

9. A platinum(II) or platinum(IV) compound according to the general formula (V),



wherein

25 n is equal to 0 or 1,


A, X are selected independently of one another from the group consisting of halogenides, OH^- , carboxylate, sulphate and sulphonate, and

Z is selected from the group consisting of hydroxyalkyl amine, hydroxyalkenyl amine, sulphoalkyl amine, sulphoalkenyl amine, which is substituted at at least one of the CH_2 - or
30 CH groups by a halogen, alkyl, alkenyl, cycloalkyl, cycloalkenyl, aryl, hydroxyl, carboxyl, sulphate,

phosphate radical and / or a heterocyclic compound and additionally the amino nitrogen can be substituted with these radicals, wherein

for n equal to 0, the two radicals Z present in the molecule can be linked via a radical selected from the group consisting of substituted or unsubstituted alkylene, alkenylene, cycloalkylene and cycloalkenylene radicals and a heterocyclic compound, preferably ethane-1,2-diyl and their physiologically acceptable addition salts.

10. A platinum(II) or platinum(IV) compound according to the general formula (VI),



wherein

A and X are selected independently of one another from the group consisting of halogenides, OH^- , OH_2 , carboxylate, sulphate and sulphonate, and

Z is selected from the group consisting of hydroxyalkyl amine, hydroxyalkenyl amine, carboxyalkyl amine, carboxyalkenyl amine, sulphoalkyl amine and sulphoalkenyl amine, which is substituted at at least one of the CH_2 - or CH groups by a halogen, alkyl, alkenyl, cycloalkyl, cycloalkenyl, aryl, hydroxyl, carboxyl, sulphate, phosphate radical and / or a heterocyclic compound and additionally can be substituted at the amino nitrogen by these radicals,

wherein

the two Z radicals present in the molecule can be linked via a radical selected from the group consisting of alkylene, alkenylene, cycloalkylene and cycloalkenylene and a heterocyclic compound, or substituted alkylene, alkenylene, cycloalkylene and cycloalkenylene


and their physiologically acceptable addition salts.

11. The compound according to claim 10, wherein two radicals Z present in the molecule are linked via ethylene.


12. The platinum(II) and / or platinum(IV) compound according to claim 9, for use as a prophylactic and / or therapeutic agent for the treatment of diseases.

13. Use of at least one platinum(II) and / or platinum(IV) compound according to claim 9, for the manufacture of a prophylactic and / or therapeutic agent for the treatment of tumor diseases.


14. Use of platinum(II) a complex of the general formulae Ib to IVb for the manufacture of medicaments for the therapy of tumor diseases,




Ib



IIb



IIIb



IVb

wherein the radicals

R^1 and $R^{3'}$ originate from the group of hydroxyalkyls and hydroxyalkenyls and hydroxyalkyls and hydroxyalkenyls substituted with halogens, alkyls, cycloalkyls, heterocyclic compounds or functional groups such as hydroxy, carboxy, sulphate or phosphate, wherein the hydroxyalkyls and hydroxyalkenyls can be present protonated or deprotonated,

$R^2, R^3, R^{1'}, R^{2'} = -CH_2-CH_2-OH, -CH_2-CH_2-CH_2-OH, -CH_2-CH_2-CH_2-CH_2-OH, -CH_2-CH_2-CH_2-CH_2-CH_2-OH, -H, \text{ methyl, ethyl, saturated or unsaturated cyclic radicals, also heterocyclic compounds, as well as their halogen, hydroxy, carboxy, sulphate or phosphate derivatives,}$

$R^4 = \text{alkyl, alkylene, cycloalkyl, cycloalkene, heterocyclic radicals or substituted alkyls and alkenes, cycloalkyl and cycloalkene, but preferably can be ethane-1,2-diyl,}$


and a belongs to the group of halogenides (fluorine, chlorine, bromine, iodine), OH , OH_2 , carboxylate, sulphate or sulphonate.

15. Use of a compound according to claim 14, wherein however R^1 and $R^{3'}$ = carboxy alkyls or carboxy alkenyls as well as carboxy alkyls and carboxy alkenyls substituted by halogens, alkyls, cycloalkyls, heterocyclic compounds or functional groups such as hydroxy, carboxy, sulphate or phosphate and the carboxy alkyls or carboxy alkenyls can be present protonated or deprotonated.

5 16. Use of a compound according to claim 14, wherein however R^1 and $R^{3'}$ = sulphoalkyls or sulphoalkenyls as well as sulphoalkyls and sulphoalkenyls substituted by halogens, alkyls, cycloalkyls, heterocyclic compounds or functional groups such as hydroxy, carboxy, sulphate or phosphate and the sulphoalkyls or sulphoalkenyls can be present protonated or deprotonated.

10 17. Use of a compound according to claim 14, wherein in particular
 R^1 and $R^{3'}$ = $-\text{CH}_2-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{COOH}$, $-\text{CH}_2-\text{CH}_2-\text{COOH}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{COOH}$, $-\text{CH}_2-\text{CH}_2-\text{SO}_3\text{H}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{SO}_3\text{H}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{SO}_3\text{H}$ and $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{SO}_3\text{H}$ and
 R^1 and $R^{3'}$ = $-\text{CH}_2-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{COOH}$, $-\text{CH}_2-\text{CH}_2-\text{COOH}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{COOH}$, $-\text{CH}_2-\text{CH}_2-\text{SO}_3\text{H}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{SO}_3\text{H}$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{SO}_3\text{H}$ and $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{SO}_3\text{H}$ can be present
15 protonated or deprotonated.

18. Use of a platinum(IV) complex with single or double intramolecular cyclizations of the general formulae Ic to IVc for the manufacture of medicaments for the therapy of tumor diseases,



wherein the radicals

R^1 and $R^{3'}$ originate from the group of hydroxyalkyls and hydroxyalkenyls, and hydroxyalkyls and hydroxyalkenyls substituted with halogens, alkyls, cycloalkyls, heterocyclic compounds or functional groups such as hydroxy, carboxy, sulphate or phosphate, wherein the
25 hydroxyalkyls and hydroxyalkenyls can be present protonated or deprotonated,

$R^2, R^3, R^1, R^2 = -CH_2-CH_2-OH, -CH_2-CH_2-CH_2-OH, -CH_2-CH_2-CH_2-CH_2-OH, -CH_2-CH_2-CH_2-CH_2-CH_2-OH, -H, \text{methyl, ethyl, saturated or unsaturated cyclic radicals, also heterocyclic compounds, as well as their halogen, hydroxy, carboxy, sulphate or phosphate derivatives,}$

5 $R^4 = \text{alkyl, alkylene, cycloalkyl, cycloalkene, heterocyclic radicals or substituted alkyls and alkenes, cycloalkyl and cycloalkene, but can preferably be ethane-1,2-diyl,}$
and a and c belong to the group of halogenides (fluorine, chlorine, bromine, iodine), $OH^-, OH_2, \text{carboxylate, sulphate or sulphonate.}$

10 19. Use of a compound according to Claim 18, wherein however R^1 and $R^{3'} = \text{carboxy alkyls or carboxy alkenyls as well as carboxy alkyls and carboxy alkenyls substituted with halogens, alkyls, cycloalkyls, heterocyclic compounds or functional groups such as hydroxy, carboxy, sulphate or phosphate and the carboxy alkyls or carboxy alkenyls can be present protonated or deprotonated.}$


15 20. Use of a compound according to claim 18, wherein however R^1 and $R^{3'} = \text{sulphoalkyls or sulphoalkenyls as well as sulphoalkyls and sulphoalkenyls substituted with halogens, alkyls, cycloalkyls, heterocyclic compounds or functional groups such as hydroxy, carboxy, sulphate or phosphate and the sulphoalkyls or sulphoalkenyls can be present protonated or deprotonated.}$

21. Use of a compound according to claim 18, in particular
20 R^1 and $R^{3'} = -CH_2-CH_2-OH, -CH_2-CH_2-CH_2-OH, -CH_2-CH_2-CH_2-CH_2-OH, -CH_2-CH_2-CH_2-CH_2-CH_2-OH, -CH_2-COOH, -CH_2-CH_2-COOH, -CH_2-CH_2-CH_2-COOH, -CH_2-CH_2-SO_3H, -CH_2-CH_2-CH_2-SO_3H, -CH_2-CH_2-CH_2-CH_2-SO_3H \text{ and } -CH_2-CH_2-CH_2-CH_2-CH_2-SO_3H \text{ and } R^1 \text{ and } R^{3'} = -CH_2-CH_2-OH, -CH_2-CH_2-CH_2-OH, -CH_2-CH_2-CH_2-CH_2-OH, -CH_2-CH_2-CH_2-CH_2-CH_2-OH, -CH_2-COOH, -CH_2-CH_2-COOH, -CH_2-CH_2-CH_2-COOH, -CH_2-CH_2-SO_3H, -CH_2-CH_2-CH_2-SO_3H, -CH_2-CH_2-CH_2-CH_2-SO_3H \text{ and } -CH_2-CH_2-CH_2-CH_2-CH_2-SO_3H \text{ can be present}$
25 protonated and deprotonated.

22. Use of a platinum compound according to claims 14, wherein however the radicals R^1 and $R^{3'}$ preferably signify substituted and non-substituted hydroxyalkyls and wherein the hydroxyalkyls can be protonated or deprotonated.

30 23. Use of a platinum compound according to claim 14, wherein however the radicals R^1 and $R^{3'}$ preferably signify 2-hydroxyethyl or deprotonated 2-hydroxyethyl, but especially preferably deprotonated 2-hydroxyethyl in the cyclized form.

24. A platinum(II) or platinum(IV) compound according to the general formula



5

wherein

n is equal to 0 or 1,

A and X belong to the group of halogenides (fluorine, chlorine, bromine, iodine),
OH⁻, OH₂, carboxylate, sulphate or sulphonate,
and Z,


10

provided n is equal to 0, the two radicals Z present in the molecule can be linked via
a radical from the group of alkyls, alkenes, cycloalkyls, cycloalkenyls, heterocyclic compounds or
substituted alkyls and alkenes, cycloalkyls and cycloalkenyls, but preferably ethane-1,2-diyl,
represents a hydroxyalkyl amine or hydroxyalkylene amine, carboxyalkyl amine or
carboxyalkylene amine or sulfoalkyl amine or sulfoalkylene amine, which is derivatized on at
least one of the CH₂ or CH groups with halogens, alkyls, cycloalkyls, heterocyclic compounds or
functional groups such as hydroxy, carboxy, sulphate or phosphate and additionally can be
derivatized on the amino nitrogen with these groups.

15

25. A platinum(II) or platinum(IV) compound according to the general formula

20



wherein

A and X belong to the group of halogenides (fluorine, chlorine, bromine, iodine),
OH⁻, OH₂, carboxylate, sulphate or sulphonate, and

25

Z represents a hydroxyalkyl amine or hydroxyalkylene amine, carboxyalkyl amine or
carboxyalkylene amine or sulfoalkyl amine or sulfoalkylene amine, which is derivatized on at
least one of the CH₂ or CH groups with halogens, alkyls, cycloalkyls, heterocyclic compounds or
functional groups such as hydroxy, carboxy, sulphate or phosphate and additionally can be
derivatized on the amino nitrogen with these groups, wherein two radicals Z present in the molecule
can be linked via a radical from the group of alkyls, alkenes, cycloalkyls, cycloalkenyls,
heterocyclic compounds or substituted alkyls and alkenes, cycloalkyls and cycloalkenyls, but
preferably ethane-1,2-diyl.

30

26. A platinum(II) or platinum(IV) compounds according to the general formula



wherein

n is equal to 0 or 1,


A and X belong to the group of halogenides (fluorine, chlorine, bromine, iodine),
OH⁻, OH₂, carboxylate, sulphate or sulphonate,
and Z,

provided n is equal to 0, the two radicals Z present in the molecule can be linked via
a radical from the group of alkyls, alkylenes, cycloalkyls, cycloalkenyls, heterocyclic compounds or
substituted alkyls and alkylenes, cycloalkyls, cycloalkenyls, but preferably ethane-1,2 diyl,

represents a hydroxyalkyl amine or hydroxyalkylene amine, carboxyalkyl amine or
carboxyalkylene amine or sulphoalkyl amine or sulphoalkylene amine, which is derivatized on at
least one of the CH₂ or CH groups with halogens, alkyls, cycloalkyls, heterocyclic compounds or
functional groups such as hydroxy, carboxy, sulphate or phosphate and additionally can be
derivatized on the amino nitrogen with these groups.

27. Use of a substance according to claim 24, for the manufacture of medicaments for the
treatment of tumor diseases.

28. Use of a platinum(II) or platinum(IV) complex of the general formula V for the
manufacture of medicaments for the therapy of tumor diseases



V

wherein the radicals R¹ and R^{3'} originate from the group of hydroxyalkyls and
hydroxyalkenyls and hydroxyalkyls and hydroxyalkenyls substituted with halogens, alkyls,

cycloalkyls, heterocyclic compounds or functional groups such as hydroxy, carboxy, sulphate or phosphate, wherein the hydroxyalkyls and hydroxyalkenyls can be present protonated or deprotonated,



- 5 $CH_2-CH_2-CH_2-CH_2-CH_2-OH$, -H, methyl, ethyl, saturated or unsaturated cyclic radicals, also heterocyclic compounds, as well as their halogen, hydroxy, carboxy, sulphate or phosphate derivatives,

R^4 = alkyl, alkylene, cycloalkyl, cycloalkene, heterocyclic radicals or substituted alkyls and alkenes, cycloalkyl and cycloalkene, but preferably ethane-1,2-diyl,

- 10 and a, a' and c, c' belong to the group of halogenides (fluorine, chlorine, bromine, iodine), OH⁻, OH₂, carboxylate, sulphate or sulphonate, wherein for platinum(II) compounds c is omitted (c0).

29. Use of a compound according to claim 28, wherein however R^1 and $R^{3'}$ = carboxy alkyls or carboxy alkenyls as well as carboxy alkyls and carboxy alkenyls substituted with halogens, alkyls, cycloalkyls, heterocyclic compounds or functional groups such as hydroxy, carboxy, sulphate or phosphate and wherein the carboxy alkyls and carboxy alkenyls can be present protonated or deprotonated.

30. Use of a compound according to claim 28, wherein however R^1 and $R^{3'}$ = sulphoalkyls or sulphoalkenyls as well as sulphoalkyls and sulphoalkenyls substituted by halogens, alkyls, cycloalkyls, heterocyclic compounds or functional groups such as hydroxy, carboxy, sulphate or phosphate and the sulphoalkyls or sulphoalkenyls can be present protonated or deprotonated.

31. Use of a compound according to claim 28, in particular

- R^1 and $R^{3'}$ = $-CH_2-CH_2-OH$, $-CH_2-CH_2-CH_2-OH$, $-CH_2-CH_2-CH_2-CH_2-OH$, $-CH_2-CH_2-CH_2-CH_2-OH$, $-CH_2-COOH$, $-CH_2-CH_2-COOH$, $-CH_2-CH_2-CH_2-COOH$, $-CH_2-CH_2-SO_3H$, $-CH_2-CH_2-CH_2-SO_3H$, $-CH_2-CH_2-CH_2-CH_2-SO_3H$ and $-CH_2-CH_2-CH_2-CH_2-CH_2-SO_3H$ and R^1 and $R^{3'}$ = $-CH_2-CH_2-OH$, $-CH_2-CH_2-CH_2-OH$, $-CH_2-CH_2-CH_2-CH_2-OH$, $-CH_2-CH_2-CH_2-CH_2-CH_2-OH$, $-CH_2-COOH$, $-CH_2-CH_2-COOH$, $-CH_2-CH_2-CH_2-COOH$, $-CH_2-CH_2-SO_3H$, $-CH_2-CH_2-CH_2-SO_3H$, $-CH_2-CH_2-CH_2-CH_2-SO_3H$ and $-CH_2-CH_2-CH_2-CH_2-CH_2-SO_3H$ can be present protonated or deprotonated.

32. Use of a platinum compound according to claim 28, wherein however the radicals R^1 and $R^{3'}$ preferably signify substituted and non-substituted hydroxyalkyls and wherein the hydroxyalkyls can be present protonated or deprotonated.

5 33. Use of a platinum compound according to claim 28, wherein however the radicals R^1 and $R^{3'}$ preferably signify 2-hydroxyethyl or deprotonated 2-hydroxyethyl, but especially preferably deprotonated 2-hydroxyethyl in the cyclized form.

34. A medicament containing a compound of claim 14.